
Cardiometabolic Food Plan—Bibliography

Scientific/Medical Publications

Why the Cardiometabolic Food Plan?

- Sperling LS, Mechanick JI, Neeland IJ, et al. The CardioMetabolic Health Alliance: working toward a new care model for the metabolic syndrome. *J Am Coll Cardiol*. 2015;66(9):1050-1067. doi:10.1016/j.jacc.2015.06.1328.
- Naka KK, Papathanassiou K, Bechlioulis A, et al. Determinants of vascular function in patients with type 2 diabetes. *Cardiovasc Diabetol*. 2012;11:127. doi:10.1186/1475-2840-11-127.
- Gordin D, Saraheimo M, Tuomikangas J, et al. Influence of postprandial hyperglycemic conditions on arterial stiffness in patients with type 2 diabetes. *J Clin Endocrinol Metab*. 2016;101(3):1134-1143. doi:10.1210/jc.2015-3635.
- Galassi A, Reynolds K, He J. Metabolic syndrome and risk of cardiovascular disease: a meta-analysis. *Am J Med*. 2006;119(10):812-819. doi:10.1016/j.amjmed.2006.02.031.
- Keys A, Menotti A, Aravanis C, et al. The seven countries study: 2,289 deaths in 15 years. *Prev Med*. 1984;13(2):141-154. doi:10.1016/0091-7435(84)90047-1.
- Trichopoulou A, Kouris-Blazos A, Wahlqvist ML, et al. Diet and overall survival in elderly people. *BMJ*. 1995;311(7018):1457-1460. doi:10.1136/bmj.311.7018.1457.
- Vardavas CI, Linardakis MK, Hatzis CM, Saris WH, Kafatos AG. Cardiovascular disease risk factors and dietary habits of farmers from Crete 45 years after the first description of the Mediterranean diet. *Eur J Cardiovasc Prev Rehabil*. 2010;17(4):440-446. doi:10.1097/HJR.0b013e32833692ea.

Features of the Cardiometabolic Food Plan

- Keys A, Menotti A, Aravanis C, et al. The seven countries study: 2,289 deaths in 15 years. *Prev Med*. 1984;13(2):141-154. doi:10.1016/0091-7435(84)90047-1.
- Trichopoulou A, Kouris-Blazos A, Wahlqvist ML, et al. Diet and overall survival in elderly people. *BMJ*. 1995;311(7018):1457-1460. doi:10.1136/bmj.311.7018.1457.

Scientific/Medical Publications

Features of the Cardiometabolic Food Plan (continued)

- Vardavas CI, Linardakis MK, Hatzis CM, Saris WH, Kafatos AG. Cardiovascular disease risk factors and dietary habits of farmers from Crete 45 years after the first description of the Mediterranean diet. *Eur J Cardiovasc Prev Rehabil*. 2010;17(4):440-446. doi:10.1097/HJR.0b013e32833692ea.
- Bonaccio M, Di Castelnuovo A, Costanzo S, Persichillo M, et al. Adherence to the traditional Mediterranean diet and mortality in subjects with diabetes. Prospective results from the MOLI-SANI study. *Eur J Prev Cardiol*. 2015 Feb 3. pii: 2047487315569409.
- Franquesa M, Pujol-Busquets G, García-Fernández E, et al. Mediterranean diet and cardiometabolic disease: a systematic review through evidence-based answers to key clinical questions. *Nutrients*. 2019;11(3):655. doi:10.3390/nu11030655.
- García-Fernández E, Rico-Cabanas L, Rosgaard N, Estruch R, et al. Mediterranean diet and cardiometabolic disease: a review. *Nutrients*. 2014 Sep 4;6(9):3474-3500.
- Rees K, Hartley L, Flowers N, Clarke A, et al. 'Mediterranean' dietary pattern for the primary prevention of cardiovascular disease. *Cochrane Database Syst Rev*. 2013 Aug 12;8:CD009825. doi: 10.1002/14651858. CD009825.pub2.
- Papadaki A, Nolen-Doerr E, Mantzoros CS. The effect of the Mediterranean diet on metabolic health: a systematic review and meta-analysis of controlled trials in adults. *Nutrients*. 2020;12(11):3342. doi:10.3390/nu12113342.
- Estruch R, Martínez-González MA, Corella D, et al. Effect of a high-fat Mediterranean diet on bodyweight and waist circumference: a prespecified secondary outcomes analysis of the PREDIMED randomised controlled trial. *Lancet Diabetes Endocrinol*. 2019;7(5):e6-e17. doi:10.1016/S2213-8587(19)30074-9
- Salas-Salvadó J, Díaz-López A, Ruiz-Canela M, et al. Effect of a lifestyle intervention program with energy-restricted Mediterranean diet and exercise on weight loss and cardiovascular risk factors: one-year results of the PREDIMED-Plus Trial. *Diabetes Care*. 2019;42(5):777-788. doi:10.2337/dc18-0836.
- Hernáez Á, Castañer O, Elosua R, et al. Mediterranean diet improves high-density lipoprotein function in high-cardiovascular-risk individuals: a randomized controlled trial. *Circulation*. 2017;135(7):633-643. doi:10.1161/CIRCULATIONAHA.116.023712.
- Davis CR, Bryan J, Hodgson JM, Woodman R, Murphy KJ. A Mediterranean diet reduces F2-isoprostanes and triglycerides among older Australian men and women after 6 months. *J Nutr*. 2017;147(7):1348-1355. doi:10.3945/jn.117.248419.

Scientific/Medical Publications

Features of the Cardiometabolic Food Plan (continued)

- Doménech M, Roman P, Lapetra J, García de la Corte FJ, et al. Mediterranean diet reduces 24-hour ambulatory blood pressure, blood glucose, and lipids: one-year randomized, clinical trial. *Hypertension*. 2014 Jul;64(1):69-76. doi: 10.1161/ HYPERTENSIONAHA.113.03353.
- Gordin D, Saraheimo M, Tuomikangas J, et al. Influence of postprandial hyperglycemic conditions on arterial stiffness in patients with type 2 diabetes. *J Clin Endocrinol Metab*. 2016;101(3):1134-1143. doi:10.1210/jc.2015-3635.
- Galassi A, Reynolds K, He J. Metabolic syndrome and risk of cardiovascular disease: a meta-analysis. *Am J Med*. 2006;119(10):812-819. doi:10.1016/j.amjmed.2006.02.031.
- Keys A, Menotti A, Aravanis C, et al. The seven countries study: 2,289 deaths in 15 years. *Prev Med*. 1984;13(2):141-154. doi:10.1016/0091-7435(84)90047-1.
- Atkinson FS, Foster-Powell K, Brand-Miller JC. International tables of glycemic index and glycemic load values: 2008. *Diabetes Care*. 2008;31(12):2281-2283. doi:10.2337/dc08-1239.
- Meng H, Matthan NR, Ausman LM, Lichtenstein AH. Effect of macronutrients and fiber on postprandial glycemic responses and meal glycemic index and glycemic load value determinations. *Am J Clin Nutr*. 2017;105(4):842-853. doi:10.3945/ajcn.116.144162.
- Sun L, Tan KWJ, Lim JZ, Magkos F, Henry CJ. Dietary fat and carbohydrate quality have independent effects on postprandial glucose and lipid responses. *Eur J Nutr*. 2018;57(1):243-250. doi:10.1007/s00394-016-1313-y.
- Turati F, Dilis V, Rossi M, Lagiou P, et al. Glycemic load and coronary heart disease in a Mediterranean population: the EPIC Greek cohort study. *Nutr Metab Cardiovasc Dis*. 2015 Mar;25(3):336-42. doi: 10.1016/j.numecd.2014.12.002.
- Chang KT, Lampe JW, Schwarz Y, Breymeyer KL, et al. Low glycemic load experimental diet more satiating than high glycemic load diet. *Nutr Cancer*. 2012;64(5):666-73. doi: 10.1080/01635581.2012.676143.
- Kraus WE, Bhapkar M, Huffman KM, et al. 2 years of calorie restriction and cardiometabolic risk (CALERIE): exploratory outcomes of a multicentre, phase 2, randomised controlled trial. *Lancet Diabetes Endocrinol*. 2019;7(9):673-683. doi:10.1016/S2213-8587(19)30151-2.
- Dashti HS, Gómez-Abellán P, Qian J, et al. Late eating is associated with cardiometabolic risk traits, obesogenic behaviors, and impaired weight loss. *Am J Clin Nutr*. 2020;nqaa264. doi:10.1093/ajcn/nqaa264.

Scientific/Medical Publications

Features of the Cardiometabolic Food Plan (continued)

- Winkvist A, Klingberg S, Nilsson LM, et al. Longitudinal 10-year changes in dietary intake and associations with cardio-metabolic risk factors in the Northern Sweden Health and Disease Study. *Nutr J.* 2017;16(1):20. doi:10.1186/s12937-017-0241-x.
- Aslani Z, Sadeghi O, Heidari-Beni M, et al. Association of dietary inflammatory potential with cardiometabolic risk factors and diseases: a systematic review and dose-response meta-analysis of observational studies. *Diabetol Metab Syndr.* 2020;12:86. doi:10.1186/s13098-020-00592-6.
- Adamska-Patrano E, Ostrowska L, Goscik J, Pietraszewska B, Kretowski A, Gorska M. The relationship between the leptin/ghrelin ratio and meals with various macronutrient contents in men with different nutritional status: a randomized crossover study. *Nutr J.* 2018;17(1):118. doi:10.1186/s12937-018-0427-x.
- Croall ID, Hoggard N, Aziz I, Hadjivassiliou M, Sanders DS. Brain fog and non-coeliac gluten sensitivity: proof of concept brain MRI pilot study. *PLoS One.* 2020;15(8):e0238283. doi:10.1371/journal.pone.0238283.
- Jiao J, Xu JY, Zhang W, Han S, Qin LQ. Effect of dietary fiber on circulating C-reactive protein in overweight and obese adults: a meta-analysis of randomized controlled trials. *Int J Food Sci Nutr.* 2015 Feb;66(1):114-9. doi:10.3109/09637486.2014.959898.
- Dahl WJ, Stewart ML. Position of the Academy of Nutrition and Dietetics: health implications of dietary fiber. *J Acad Nutr Diet.* 2015;115(11):1861-1870. doi:10.1016/j.jand.2015.09.003.
- Reynolds A, Mann J, Cummings J, Winter N, Mete E, Te Morenga L. Carbohydrate quality and human health: a series of systematic reviews and meta-analyses [published correction appears in *Lancet.* 2019 Feb 2;393(10170):406]. *Lancet.* 2019;393(10170):434-445. doi:10.1016/S0140-6736(18)31809-9.
- Delargy HJ, O'Sullivan KR, Fletcher RJ, Blundell JE. Effects of amount and type of dietary fibre (soluble and insoluble) on short-term control of appetite. *Int J Food Sci Nutr.* 1997;48(1):67-77. doi:10.3109/09637489709006965.
- Chandalia M, Garg A, Lutjohann D, von Bergmann K, Grundy SM, Brinkley LJ. Beneficial effects of high dietary fiber intake in patients with type 2 diabetes mellitus. *N Engl J Med.* 2000;342(19):1392-1398. doi:10.1056/NEJM200005113421903.
- Rees K, Dyakova M, Wilson N, Ward K, et al. Dietary advice for reducing cardiovascular risk. *Cochrane Database Syst Rev.* 2013 Dec 6;12:CD002128. doi:10.1002/14651858.CD002128.pub5.

Scientific/Medical Publications

Features of the Cardiometabolic Food Plan (continued)

- Chen GC, Lv DB, Pang Z, Dong JY, Liu QF. Dietary fiber intake and stroke risk: a meta-analysis of prospective cohort studies. *Eur J Clin Nutr.* 2013 Jan;67(1):96-100. doi: 10.1038/ejcn.2012.158.
- Gaesser GA, Rodriguez J, Patrie JT, Whisner CM, Angadi SS. Effects of glyce-mic index and cereal fiber on postprandial endothelial function, glycemia, and insulinemia in healthy adults. *Nutrients.* 2019;11(10):2387. doi:10.3390/nu1102387.
- Wu Y, Qian Y, Pan Y, Li P, et al. Association between dietary fiber intake and risk of coronary heart disease: A meta-analysis. *Clin Nutr.* 2015 Aug;34(4):603-11. doi: 10.1016/j.clnu.2014.05.009.
- Stanhope KL, Medici V, Bremer AA, Lee V, et al. A dose-response study of consuming high-fructose corn syrup- sweetened beverages on lipid/lipoprotein risk factors for cardiovascular disease in young adults. *Am J Clin Nutr.* 2015 Jun;101(6):1144-54. doi: 10.3945/ajcn.114.100461.
- Wang J. Consumption of added sugars and development of metabolic syn-drome components among a sample of youth at risk of obesity. *Appl Physiol Nutr Metab.* 2014 Apr;39(4):512. doi: 10.1139/apnm-2013-0456.
- Xi B, Huang Y, Reilly KH, et al. Sugar-sweetened beverages and risk of hyperten-sion and CVD: a dose-response meta-analysis. *Br J Nutr.* 2015;113(5):709-717. doi:10.1017/S0007114514004383.
- Qin P, Li Q, Zhao Y, et al. Sugar and artificially sweetened beverages and risk of obesity, type 2 diabetes mellitus, hypertension, and all-cause mortality: a dose-response meta-analysis of prospective cohort studies. *Eur J Epidemiol.* 2020;35(7):655-671. doi:10.1007/s10654-020-00655-y
- Raatz SK, Johnson LK, Picklo MJ. Consumption of honey, sucrose, and high-fructose corn syrup produces similar metabolic effects in glucose-tolerant and -intolerant individuals. *J Nutr.* 2015;145(10):2265-2272. doi:10.3945/jn.115.218016.
- Imamura F, O'Connor L, Ye Z, et al. Consumption of sugar sweetened beverages, artificially sweetened beverages, and fruit juice and incidence of type 2 diabetes: systematic review, meta-analysis, and estimation of population attributable fraction. *BMJ.* 2015;351:h3576. doi:10.1136/bmj.h3576.
- Sadeghi F, Akhlaghi M, Salehi S. Adverse effects of honey on low-density lipo-protein cholesterol and adiponectin concentrations in patients with type 2 diabetes: a randomized controlled cross-over trial. *J Diabetes Metab Disord.* 2020;19(1):373-380. doi:10.1007/s40200-020-00518-z.

Scientific/Medical Publications

Features of the Cardiometabolic Food Plan (continued)

- Luger M, Lafontan M, Bes-Rastrollo M, Winzer E, Yumuk V, Farpour-Lambert N. Sugar-sweetened beverages and weight gain in children and adults: a systematic review from 2013 to 2015 and a comparison with previous studies. *Obes Facts*. 2017;10(6):674-693. doi:10.1159/000484566.
- Cochrane Review includes SSB & Weight Gain von Philipsborn P, Stratil JM, Burns J, et al. Environmental interventions to reduce the consumption of sugar-sweetened beverages and their effects on health. *Cochrane Database Syst Rev*. 2019;6(6):CD012292. doi:10.1002/14651858.CD012292.pub2.
- Van Name M, Giannini C, Santoro N, et al. Blunted suppression of acyl-ghrelin in response to fructose ingestion in obese adolescents: the role of insulin resistance. *Obesity (Silver Spring)*. 2015;23(3):653-661. doi:10.1002/oby.21019.
- Guasch-Ferré M, Babio N, Martínez-González MA, et al. Dietary fat intake and risk of cardiovascular disease and all-cause mortality in a population at high risk of cardiovascular disease. *Am J Clin Nutr*. 2015;102(6):1563-1573. doi:10.3945/ajcn.115.116046.
- Luukkonen PK, Sädevirta S, Zhou Y, et al. Saturated fat is more metabolically harmful for the human liver than unsaturated fat or simple sugars. *Diabetes Care*. 2018;41(8):1732-1739. doi:10.2337/dc18-0071.
- Liu Q, Matthan NR, Manson JE, et al. Plasma phospholipid fatty acids and coronary heart disease risk: a matched case-control study within the Women's Health Initiative Observational Study. *Nutrients*. 2019;11(7):1672. doi:10.3390/nu11071672.
- Zong G, Li Y, Wanders AJ, et al. Intake of individual saturated fatty acids and risk of coronary heart disease in US men and women: two prospective longitudinal cohort studies. *BMJ*. 2016;355:i5796. doi:10.1136/bmj.i5796.
- Berrougui H, Ikhlef S, Khalil A. Extra virgin olive oil polyphenols promote cholesterol efflux and improve HDL functionality. *Evid Based Complement Alternat Med*. 2015;2015:208062. doi:10.1155/2015/208062.
- Abdelhamid AS, Brown TJ, Brainard JS, et al. Omega-3 fatty acids for the primary and secondary prevention of cardiovascular disease. *Cochrane Database Syst Rev*. 2018;7(7):CD003177. doi:10.1002/14651858.CD003177.pub3.
- Liou YA, King DJ, Zibrik D, Innis SM. Decreasing linoleic acid with constant alpha-linolenic acid in dietary fats increases (n-3) eicosapentaenoic acid in plasma phospholipids in healthy men. *J Nutr*. 2007;137(4):945-952. doi:10.1093/jn/137.4.945.
- Zhou D, Yu H, He F, Reilly KH, et al. Nut consumption in relation to cardiovascular disease risk and type 2 diabetes: a systematic review and meta-analysis of prospective studies. *Am J Clin Nutr*. 2014 May 7;100(1):270-277.

Scientific/Medical Publications

Features of the Cardiometabolic Food Plan (continued)

- Egert S, Baxheinrich A, Lee-Barkey YH, Tschoepe D, et al. Effects of an energy-restricted diet rich in plant-derived alpha-linolenic acid on systemic inflammation and endothelial function in overweight-to-obese patients with metabolic syndrome traits. *Br J Nutr.* 2014 Oct 28;112(8):1315-22. doi: 10.1017/S0007114514002001.
- Bosch J, Gerstein HC, et al. n-3 fatty acids and cardiovascular outcomes in patients with dysglycemia. *N Engl J Med.* 2012;367(4):309-318. doi:10.1056/NEJMOa1203859.
- Fraga CG, Croft KD, Kennedy DO, Tomás-Barberán FA. The effects of polyphenols and other bioactives on human health. *Food Funct.* 2019;10(2):514-528. doi:10.1039/c8fo01997e.
- Murphy MM, Barraj LM, Spungen JH, Herman DR, Randolph RK. Global assessment of select phytonutrient intakes by level of fruit and vegetable consumption. *Br J Nutr.* 2014;112(6):1004-1018. doi:10.1017/S0007114514001937.
- Castro-Acosta ML, Smith L, Miller RJ, McCarthy DI, Farrimond JA, Hall WL. Drinks containing anthocyanin-rich blackcurrant extract decrease postprandial blood glucose, insulin and incretin concentrations. *J Nutr Biochem.* 2016;38:154-161. doi:10.1016/j.jnutbio.2016.09.002.
- Hausenblas HA, Schoulda JA, Smoliga JM. Resveratrol treatment as an adjunct to pharmacological management in type 2 diabetes mellitus--systematic review and meta-analysis. *Mol Nutr Food Res.* 2015;59(1):147-159. doi:10.1002/mnfr.201400173.
- Wang Z, Huang Y, Zou J, Cao K, Xu Y, Wu JM. Effects of red wine and wine polyphenol resveratrol on platelet aggregation in vivo and in vitro. *Int J Mol Med.* 2002;9(1):77-79.
- Chiva-Blanch G, Badimon L. Effects of polyphenol intake on metabolic syndrome: current evidences from human trials. *Oxid Med Cell Longev.* 2017;2017:5812401. doi:10.1155/2017/5812401.
- Fuangchan A, Sonthisombat P, Seubnukarn T, et al. Hypoglycemic effect of bitter melon compared with metformin in newly diagnosed type 2 diabetes patients. *J Ethnopharmacol.* 2011;134(2):422-428. doi:10.1016/j.jep.2010.12.045.
- Yin RV, Lee NC, Hirpara H, Phung OJ. The effect of bitter melon (*Mormordica charantia*) in patients with diabetes mellitus: a systematic review and meta-analysis. *Nutr Diabetes.* 2014;4(12):e145. doi:10.1038/nutd.2014.42.

Scientific/Medical Publications

Features of the Cardiometabolic Food Plan (continued)

- Brüll V, Burak C, Stoffel-Wagner B, et al. Effects of a quercetin-rich onion skin extract on 24 h ambulatory blood pressure and endothelial function in overweight-to-obese patients with (pre-)hypertension: a randomised double-blinded placebo-controlled cross-over trial. *Br J Nutr.* 2015;114(8):1263-1277. doi:10.1017/S0007114515002950.
- Komorita Y, Iwase M, Fujii H, et al. Additive effects of green tea and coffee on all-cause mortality in patients with type 2 diabetes mellitus: the Fukuoka Diabetes Registry. *BMJ Open Diabetes Res Care.* 2020;8(1):e001252. doi:10.1136/bmjdr-2020-001252.
- Kouzi SA, Yang S, Nuzum DS, Dirks-Naylor AJ. Natural supplements for improving insulin sensitivity and glucose uptake in skeletal muscle. *Front Biosci (Elite Ed).* 2015;7:94-106. Published 2015 Jan 1.
- Allen RW, Schwartzman E, Baker WL, Coleman CI, Phung OJ. Cinnamon use in type 2 diabetes: an updated systematic review and meta-analysis. *Ann Fam Med.* 2013;11(5):452-459. doi:10.1370/afm.1517.
- Ranasinghe P, Jayawardena R, Pigera S, et al. Evaluation of pharmacodynamic properties and safety of *Cinnamomum zeylanicum* (Ceylon cinnamon) in healthy adults: a phase I clinical trial. *BMC Complement Altern Med.* 2017;17(1):550. doi:10.1186/s12906-017-2067-7.
- Langella C, Naviglio D, Marino M, Gallo M. Study of the effects of a diet supplemented with active components on lipid and glycemic profiles. *Nutrition.* 2015 Jan;31(1):180-6. doi: 10.1016/j.nut.2014.07.015.
- Ho HV, Sievenpiper JL, Zurbau A, et al. A systematic review and meta-analysis of randomized controlled trials of the effect of barley B-glucan on LDL-C, non-HDL-C and apoB for cardiovascular disease risk reduction. *Eur J Clin Nutr.* 2016;70(11):1239-1245. doi:10.1038/ejcn.2016.89.
- Aune D, Keum N, Giovannucci E, et al. Dietary intake and blood concentrations of antioxidants and the risk of cardiovascular disease, total cancer, and all-cause mortality: a systematic review and dose-response meta-analysis of prospective studies. *Am J Clin Nutr.* 2018;108(5):1069-1091. doi:10.1093/ajcn/nqy097.
- Fang K, Dong H, Wang D, Gong J, Huang W, Lu F. Soy isoflavones and glucose metabolism in menopausal women: A systematic review and meta-analysis of randomized controlled trials. *Mol Nutr Food Res.* 2016;60(7):1602-1614. doi:10.1002/mnfr.201501024.
- Almario RU, Karakas SE. Lignan content of the flaxseed influences its biological effects in healthy men and women. *J Am Coll Nutr.* 2013;32(3):194-9. doi: 10.1080/07315724.2013.791147.

Scientific/Medical Publications

Features of the Cardiometabolic Food Plan (continued)

- Godos J, Vitale M, Micek A, et al. Dietary polyphenol intake, blood pressure, and hypertension: a systematic review and meta-analysis of observational studies. *Antioxidants (Basel)*. 2019;8(6):152. doi:10.3390/antiox8060152.
- Johnson SA, Figueroa A, Navaei N, Wong A, et al. Daily blueberry consumption improves blood pressure and arterial stiffness in postmenopausal women with pre- and stage 1-hypertension: a randomized, double-blind, placebo-controlled clinical trial. *J Acad Nutr Diet*. 2015 Mar;115(3):369-77. doi: 10.1016/j.jand.2014.11.001.

Touring the Cardiometabolic Food Plan

- Meng H, Matthan NR, Ausman LM, Lichtenstein AH. Effect of macronutrients and fiber on postprandial glycemic responses and meal glycemic index and glycemic load value determinations. *Am J Clin Nutr*. 2017;105(4):842-853. doi:10.3945/ajcn.116.144162.
- Pasiakos SM, Lieberman HR, Fulgoni VL. Higher-protein diets are associated with higher HDL cholesterol and lower BMI and waist circumference in US adults. *J Nutr*. 2015 Mar;145(3):605-14. doi: 10.3945/jn.114.205203.
- Downer MK, Martínez-González MA, Gea A, et al. Mercury exposure and risk of cardiovascular disease: a nested case-control study in the PREDIMED (PREvention with MEDiterranean Diet) study. *BMC Cardiovasc Disord*. 2017;17(1):9. doi:10.1186/s12872-016-0435-8.
- Sluijs I, Plantinga Y, de Roos B, Mennen LI, Bots ML. Dietary supplementation with cis-9,trans-11 conjugated linoleic acid and aortic stiffness in overweight and obese adults. *Am J Clin Nutr*. 2010;91(1):175-183. doi:10.3945/ajcn.2009.28192.
- Bakhtiary A, Yassin Z, Hanachi P, Rahmat A, et al. Effects of soy on metabolic biomarkers of cardiovascular disease in elderly women with metabolic syndrome. *Arch Iran Med*. 2012 Aug;15(8):462-8. doi: 012158/AIM.004.
- Langella C, Naviglio D, Marino M, Gallo M. Study of the effects of a diet supplemented with active components on lipid and glycemic profiles. *Nutrition*. 2015 Jan;31(1):180-6. doi: 10.1016/j.nut.2014.07.015.
- Cugnet-Anceau C, Nazare JA, Biorklund M, et al. A controlled study of consumption of beta-glucan-enriched soups for 2 months by type 2 diabetic free-living subjects. *Br J Nutr*. 2010;103(3):422-428. doi:10.1017/S0007114509991875.
- Ho HV, Sievenpiper JL, Zurbau A, et al. A systematic review and meta-analysis of randomized controlled trials of the effect of barley B-glucan on LDL-C, non-HDL-C and apoB for cardiovascular disease risk reductions. 2016. *Eur J Clin Nutr*. 2016;70(11):1239-1245. doi:10.1038/ejcn.2016.89.

Scientific/Medical Publications

Touring the Cardiometabolic Food Plan (continued)

- Ho HV, Sievenpiper JL, Zurbau A, Blanco Mejia S, Jovanovski E, Au-Yeung F, Jenkins AL, Vuksan V. The effect of oat B-glucan on LDL-cholesterol, non-HDL-cholesterol and apoB for CVD risk reduction: a systematic review and meta-analysis of randomised-controlled trials. *Br J Nutr.* 2016 Oct;116(8):1369-1382. doi: 10.1017/S000711451600341X.
- Widmer RJ, Freund MA, Flammer AJ, et al. Beneficial effects of polyphenol-rich olive oil in patients with early atherosclerosis. *Eur J Nutr.* 2013;52(3):1223-1231. doi:10.1007/s00394-012-0433-2.
- Medina-Remón A, Tresserra-Rimbau A, Pons A, Tur JA, et al. Effects of total dietary polyphenols on plasma nitric oxide and blood pressure in a high cardiovascular risk cohort. The PREDIMED randomized trial. *Nutr Metab Cardiovasc Dis.* 2015 Jan;25(1):60-7. doi: 10.1016/j.numecd.2014.09.001.
- Gorzynik-Debicka M, Przychodzen P, Cappello F, et al. Potential health benefits of olive oil and plant polyphenols. *Int J Mol Sci.* 2018;19(3):686. doi:10.3390/ijms19030686.
- Colica C, Di Renzo L, Trombetta D, et al. Antioxidant effects of a hydroxytyrosol-based pharmaceutical formulation on body composition, metabolic state, and gene expression: a randomized double-blinded, placebo-controlled crossover trial. *Oxid Med Cell Longev.* 2017;2017:2473495. doi:10.1155/2017/2473495.
- Perrone MA, Gualtieri P, Gratteri S, et al. Effects of postprandial hydroxytyrosol and derivatives on oxidation of LDL, cardiometabolic state and gene expression: a nutrigenomic approach for cardiovascular prevention. *J Cardiovasc Med (Hagerstown).* 2019;20(7):419-426. doi:10.2459/JCM.0000000000000816.
- Austria JA, Richard MN, Chahine MN, et al. Bioavailability of alpha-linolenic acid in subjects after ingestion of three different forms of flaxseed. *J Am Coll Nutr.* 2008;27(2):214-221.
- Patenaude A, Rodriguez-Leyva D, Edel AL, et al. Bioavailability of alpha-linolenic acid from flaxseed diets as a function of the age of the subject. *Eur J Clin Nutr.* 2009;63(9):1123-1129. doi:10.1038/ejcn.2009.41.
- Almario RU, Karakas SE. Lignan content of the flaxseed influences its biological effects in healthy men and women. *J Am Coll Nutr.* 2013;32(3):194-9. doi: 10.1080/07315724.2013.791147.
- Saleh-Ghadimi S, Kheirouri S, Golmohammadi A, Moludi J, Jafari-Vayghan H, Alizadeh M. Effect of flaxseed oil supplementation on anthropometric and metabolic indices in patients with coronary artery disease: a double-blinded randomized controlled trial. *J Cardiovasc Thorac Res.* 2019;11(2):152-160. doi:10.15171/jcvtr.2019.26.

Scientific/Medical Publications

Touring the Cardiometabolic Food Plan (continued)

- Haghighatsiar N, Askari G, Saraf-Bank S, Feizi A, Keshmiri H. Effect of flaxseed powder on cardiovascular risk factor in dyslipidemic and hypertensive patients. *Int J Prev Med*. 2019;10:218. doi:10.4103/ijpvm.IJPVM_563_17.
- Yari Z, Rahimlou M, Poustchi H, Hekmatdoost A. Flaxseed supplementation in metabolic syndrome management: a pilot randomized, open-labeled, controlled study. *Phytother Res*. 2016;30(8):1339-1344. doi:10.1002/ptr.5635.
- Liu G, Guasch-Ferré M, Hu Y, et al. Nut consumption in relation to cardiovascular disease incidence and mortality among patients with diabetes mellitus. *Circ Res*. 2019;124(6):920-929. doi:10.1161/CIRCRESAHA.118.314316.
- Zhou D, Yu H, He F, Reilly KH, et al. Nut consumption in relation to cardiovascular disease risk and type 2 diabetes: a systematic review and meta-analysis of prospective studies. *Am J Clin Nutr*. 2014 May 7;100(1):270-277.
- Wannamethee SG, Jefferis BJ, Lennon L, Papacosta O, Whincup PH, Hingorani AD. Serum conjugated linoleic acid and risk of incident heart failure in older men: The British Regional Heart Study. *J Am Heart Assoc*. 2018;7(1):e006653. doi:10.1161/JAHA.117.006653.
- Bakhtiary A, Yassin Z, Hanachi P, Rahmat A, et al. Effects of soy on metabolic biomarkers of cardiovascular disease in elderly women with metabolic syndrome. *Arch Iran Med*. 2012 Aug;15(8):462-8. doi: 012158/AIM.004.
- van Nielen M, Feskens EJ, Rietman A, Siebelink E, et al. Partly replacing meat protein with soy protein alters insulin resistance and blood lipids in postmenopausal women with abdominal obesity. *J Nutr*. 2014 Sep;144(9):1423-9. doi: 10.3945/jn.114.193706.
- Masley SC, Roetzheim R, Masley LV, McNamara T, Schocken DD. Emerging risk factors as markers for carotid intima media thickness scores. *J Am Coll Nutr*. 2015 Mar-Apr;34(2) 100-107. doi:10.1080/07315724.2014.916238.
- Man, B. Cui, C, Zhang, X, et al. The effect of soy isoflavones on arterial stiffness: a systematic review and meta-analysis of randomized controlled trials. *Eur J Nutr* (2020). doi.org/10.1007/s00394-020-02300-6.
- Lin YJ, Lin P. Probabilistic integrated human mixture risk assessment of multiple metals through seafood consumption. *Risk Anal*. 2019;39(2):426-438. doi:10.1111/risa.13183.
- Bondonno CP, Blekkenhorst LC, Prince RL, et al. Association of vegetable nitrate intake with carotid atherosclerosis and ischemic cerebrovascular disease in older women. *Stroke*. 2017;48(7):1724-1729. doi:10.1161/strokeaha.117.016844.
- Brkic D, Bošnjir J, Bevardi M, et al. Nitrate in leafy green vegetables and estimated intake. *Afr J Tradit Complement Altern Med*. 2017;14(3):31-41. doi:10.21010/ajtcam.v14i3.4.

Scientific/Medical Publications

Touring the Cardiometabolic Food Plan (continued)

- Bazzano LA, He J, Ogden LG, et al. Dietary intake of folate and risk of stroke in US men and women: NHANES I Epidemiologic Follow-up Study. National Health and Nutrition Examination Survey. *Stroke*. 2002;33(5):1183-1188. doi:10.1161/01.str.0000014607.90464.88.
- Song B, Liu K, Gao Y, et al. Lycopene and risk of cardiovascular diseases: a meta-analysis of observational studies. *Mol Nutr Food Res*. 2017;61(9):10.1002/mnfr.201601009. doi:10.1002/mnfr.201601009.
- Cheng HM, Koutsidis G, Lodge JK, Ashor A, Siervo M, Lara J. Tomato and lycopene supplementation and cardiovascular risk factors: a systematic review and meta-analysis. *Atherosclerosis*. 2017;257:100-108. doi:10.1016/j.atherosclerosis.2017.01.009.
- Gadelha CJMU, Bezerra AN. Effects of probiotics on the lipid profile: systematic review. *J Vasc Bras*. 2019;18:e20180124. doi:10.1590/1677-5449.180124.
- Cassidy A, Bertoia M, Chiuve S, Flint A, Forman J, Rimm EB. Habitual intake of anthocyanins and flavanones and risk of cardiovascular disease in men. *Am J Clin Nutr*. 2016;104(3):587-594. doi:10.3945/ajcn.116.133132.
- Curtis PJ, van der Velpen V, Berends L, et al. Blueberries improve biomarkers of cardiometabolic function in participants with metabolic syndrome-results from a 6-month, double-blind, randomized controlled trial [published correction appears in *Am J Clin Nutr*. 2019 Nov 1;110(5):1262]. *Am J Clin Nutr*. 2019;109(6):1535-1545. doi:10.1093/ajcn/nqy380.
- Stull AJ, Cash KC, Champagne CM, et al. Blueberries improve endothelial function, but not blood pressure, in adults with metabolic syndrome: a randomized, double-blind, placebo-controlled clinical trial. *Nutrients*. 2015;7(6):4107-4123. doi:10.3390/nu7064107.
- Sohrab G, Roshan H, Ebrahimof S, Nikpayam O, Sotoudeh G, Siasi F. Effects of pomegranate juice consumption on blood pressure and lipid profile in patients with type 2 diabetes: a single-blind randomized clinical trial. *Clin Nutr ESPEN*. 2019;29:30-35. doi:10.1016/j.clnesp.2018.11.013.
- Mirmiran P, Fazeli MR, Asghari G, Shafiee A, Azizi F. Effect of pomegranate seed oil on hyperlipidaemic subjects: a double-blind placebo-controlled clinical trial. *Br J Nutr*. 2010;104(3):402-406. doi:10.1017/S0007114510000504.
- Li X, Cai X, Ma X, et al. Short- and Long-Term Effects of wholegrain oat intake on weight management and glucolipid metabolism in overweight type-2 diabetics: a randomized control trial. *Nutrients*. 2016;8(9):549. doi:10.3390/nu8090549.
- Bao L, Cai X, Xu M, Li Y. Effect of oat intake on glycaemic control and insulin sensitivity: a meta-analysis of randomised controlled trials. *Br J Nutr*. 2014;112(3):457-466. doi:10.1017/S0007114514000889.

Scientific/Medical Publications

Touring the Cardiometabolic Food Plan (continued)

- Baba S, Natsume M, Yasuda A, et al. Plasma LDL and HDL cholesterol and oxidized LDL concentrations are altered in normo- and hypercholesterolemic humans after intake of different levels of cocoa powder. *J Nutr.* 2007;137(6):1436-1441. doi:10.1093/jn/137.6.1436.
- Jia L, Liu X, Bai YY, et al. Short-term effect of cocoa product consumption on lipid profile: a meta-analysis of randomized controlled trials. *Am J Clin Nutr.* 2010;92(1):218-225. doi:10.3945/ajcn.2009.28202.
- Komorita Y, Iwase M, Fujii H, et al. Additive effects of green tea and coffee on all-cause mortality in patients with type 2 diabetes mellitus: the Fukuoka Diabetes Registry. *BMJ Open Diabetes Res Care.* 2020;8(1):e001252. doi:10.1136/bmjdr-2020-001252.
- Wells BJ, Mainous AG 3rd, Everett CJ. Association between dietary arginine and C-reactive protein. *Nutrition.* 2005;21(2):125-130. doi:10.1016/j.nut.2004.03.021.
- Eady S, Wallace A, Willis J, Scott R, Frampton C. Consumption of a plant sterol-based spread derived from rice bran oil is effective at reducing plasma lipid levels in mildly hypercholesterolaemic individuals. *Br J Nutr.* 2011;105(12):1808-1818. doi:10.1017/S0007114510005519.

Frequently Asked Questions

- Chagas P, Caramori P, Galdino TP, Barcellos Cda S, et al. Egg consumption and coronary atherosclerotic burden. *Atherosclerosis.* 2013 Aug;229(2):381-4. doi:10.1016/j.atherosclerosis.2013.05.008.
- Hausenblas HA, Schoulda JA, Smoliga JM. Resveratrol treatment as an adjunct to pharmacological management in type 2 diabetes mellitus--systematic review and meta-analysis. *Mol Nutr Food Res.* 2015;59(1):147-159. doi:10.1002/mnfr.201400173.
- Ma H, Malone KE, McDonald JA, et al. Pre-diagnosis alcohol consumption and mortality risk among black women and white women with invasive breast cancer. *BMC Cancer.* 2019;19(1):800. doi:10.1186/s12885-019-5991-8.
- Janssen I, Landay AL, Ruppert K, Powell LH. Moderate wine consumption is associated with lower hemostatic and inflammatory risk factors over 8 years: the study of women's health across the nation (SWAN). *Nutr Aging (Amst).* 2014 Jun 12;2(2-3):91-99.

