


Effects of Physical Inactivity

Your body was designed to move. If you don't exercise enough and sit too much, you are at greater risk of chronic diseases, especially as you age. These chronic health concerns include heart disease, type 2 diabetes, certain cancers, osteoporosis, dementia, and depression.^{1,2} The following diagram shows potential harmful changes in the body and health risks linked with sitting too much and not getting enough physical activity.



Guidelines for Physical Activity

Adults need both aerobic (cardio) and muscle-strengthening activity to support health.¹ Examples of aerobic activity are brisk walking, jogging, dancing, bicycling, and swimming. Examples of muscle-strengthening activity are lifting weights, using resistance bands, and doing body-weight exercises like push-ups. Finding a variety of activities you enjoy is key. Below are guidelines for each activity type.

Aerobic Activity

Aim for 150 to 300 minutes of moderate-intensity aerobic activity or 75 to 150 minutes of vigorous-intensity aerobic activity per week. If desired, you can do a mix of moderate- and vigorous-intensity aerobic activity. In general, 1 minute of vigorous-intensity activity equals 2 minutes of moderate-intensity activity.¹

- Gauge intensity with the talk test: If you can talk but not sing during physical activity, it is likely moderate-intensity activity. If you can't say more than a few words without pausing for a breath, it is likely vigorous-intensity activity.¹
- High-intensity interval training (HIIT), which alternates between short bursts of intense exercise and brief recovery periods, is considered vigorous-intensity activity.¹
- Spread your aerobic activity minutes throughout the week, preferably across three or more days.¹
- If you are already pretty active, consider progressing to more than 300 minutes of moderate-intensity activity per week.¹ However, avoid overexercise, which is signaled by an inability to recover between workouts or declining performance.³⁴

Muscle Strengthening Activity

Do muscle-strengthening activity of moderate or greater intensity at least two days a week.¹

- Repeat strength exercises until the muscle is fatigued. For example, you might do 8 to 12 repetitions of bicep curls.
- Work all major muscle groups of your body, including the arms, shoulders, legs, hips, back, chest, and abdomen.
- Avoid exercising the same muscle groups two days in a row. Your muscles need time to recover.³⁵

Sitting Less

Getting the recommended amount of physical activity per week is not enough to avoid all of the health risks of excess sitting.³⁶ Make a special effort to sit less at work and home. Try to incorporate easy exercises, standing breaks, or create active workstations³⁷ to help break up long periods of sitting.

Readiness for Physical Activity

Remember, if you have a chronic health condition, such as diabetes or heart disease, consult your primary care physician before starting a physical activity program.¹ Some other tips to keep in mind include:

- Choose activities suited to your current fitness level and abilities.
- Consult a physical activity specialist for guidance on appropriate activities.
- Do your best. If your current fitness level or a chronic condition prevents you from doing the minimum 150 minutes a week of moderate-intensity activity, start slow. Even adding small amounts of physical activity can support health.¹

REFERENCES

1. U.S. Department of Health and Human Services. Physical Activity Guidelines for Americans. 2nd ed. Washington, DC: U.S. Department of Health and Human Services; 2018. Accessed May 28, 2025. https://odphp.health.gov/sites/default/files/2019-09/Physical_Activity_Guidelines_2nd_edition.pdf
2. Park JH, Moon JH, Kim HJ, Kong MH, Oh YH. Sedentary lifestyle: overview of updated evidence of potential health risks. *Korean J Fam Med*. 2020;41(6):365-373. doi:10.4082/kjfm.20.0165
3. Huang Y, Li L, Gan Y, et al. Sedentary behaviors and risk of depression: a meta-analysis of prospective studies. *Transl Psychiatry*. 2020;10:26. doi:10.1038/s41398-020-0715-z
4. Wheeler MJ, Green DJ, Ellis KA, et al. Distinct effects of acute exercise and breaks in sitting on working memory and executive function in older adults: a three-arm, randomised cross-over trial to evaluate the effects of exercise with and without breaks in sitting on cognition. *Br J Sports Med*. 2020;54(13):776-781. doi:10.1136/bjsports-2018-100168
5. Tian S, Liang Z, Qiu F, Wang X. Physical activity on executive function in sedentary individuals: Systematic review and meta-analysis of randomized controlled trials. *PLOS ONE*. 2023;18(12):e0294251. doi:10.1371/journal.pone.0294251
6. Zou L, Herold F, Cheval B, et al. Sedentary behavior and lifespan brain health. *Trends Cogn Sci*. 2024;28(4):369-382. doi:10.1016/j.tics.2024.02.003
7. Sun Y, Chen C, Yu Y, et al. Replacement of leisure-time sedentary behavior with various physical activities and the risk of dementia incidence and mortality: A prospective cohort study. *J Sport Health Sci*. 2023;12(3):287-294. doi:10.1016/j.jshs.2022.11.005
8. Raichlen DA, Aslan DH, Sayre MK, et al. Sedentary behavior and incident dementia among older adults. *JAMA*. 2023;330(10):934-940. doi:10.1001/jama.2023.15231
9. Smart NA, King N, McFarlane JR, Graham PL, Dieberg G. Effect of exercise training on liver function in adults who are overweight or exhibit fatty liver disease: a systematic review and meta-analysis. *Br J Sports Med*. 2018;52(13):834-843. doi:10.1136/bjsports-2016-096197
10. Kuan WH, Chen YL, Liu CL. Excretion of Ni, Pb, Cu, As, and Hg in sweat under two sweating conditions. *Int J Environ Res Public Health*. 2022;19(7):4323. doi:10.3390/ijerph19074323
11. Wu J, Zhang H, Yang L, et al. Sedentary time and the risk of metabolic syndrome: A systematic review and dose-response meta-analysis. *Obes Rev Off J Int Assoc Study Obes*. 2022;23(12):e13510. doi:10.1111/obr.13510
12. Quan M, Xun P, Wu H, et al. Effects of interrupting prolonged sitting on postprandial glycemia and insulin responses: A network meta-analysis. *J Sport Health Sci*. 2021;10(4):419-429. doi:10.1016/j.jshs.2020.12.006
13. Hadgraft NT, Winkler E, Climie RE, et al. Effects of sedentary behaviour interventions on biomarkers of cardiometabolic risk in adults: systematic review with meta-analyses. *Br J Sports Med*. 2021;55(3):144-154. doi:10.1136/bjsports-2019-101154
14. Gubelmann C, Kuehner C, Vollenweider P, Marques-Vidal P. Association of activity status and patterns with salivary cortisol: the population-based CoLaus study. *Eur J Appl Physiol*. 2018;118(7):1507-1514. doi:10.1007/s00421-018-3881-4
15. Chantry AJ, Bishop NC, Hamer M, Kingsnorth AP, Chen YL, Paine NJ. Sedentary behaviour is associated with heightened cardiovascular, inflammatory and cortisol reactivity to acute psychological stress. *Psychoneuroendocrinology*. 2022;141:105756. doi:10.1016/j.psyneuen.2022.105756
16. Hermelink R, Leitzmann MF, Markozannes G, et al. Sedentary behavior and cancer—an umbrella review and meta-analysis. *Eur J Epidemiol*. 2022;37(5):447-460. doi:10.1007/s10654-022-00873-6
17. Friedenreich CM, Ryder-Burbidge C, McNeil J. Physical activity, obesity and sedentary behavior in cancer etiology: epidemiologic evidence and biologic mechanisms. *Mol Oncol*. 2021;15(3):790-800. doi:10.1002/1878-0261.12772
18. Izquierdo M, Merchant RA, Morley JE, et al. International exercise recommendations in older adults (ICFSR): Expert Consensus Guidelines. *J Nutr Health Aging*. 2021;25(7):824-853. doi:10.1007/s12603-021-1665-8
19. Shur NF, Creedon L, Skirrow S, et al. Age-related changes in muscle architecture and metabolism in humans: The likely contribution of physical inactivity to age-related functional decline. *Ageing Res Rev*. 2021;68:101344. doi:10.1016/j.arr.2021.101344
20. Pinheiro MB, Oliveira J, Bauman A, Fairhall N, Kwok W, Sherrington C. Evidence on physical activity and osteoporosis prevention for people aged 65+ years: a systematic review to inform the WHO guidelines on physical activity and sedentary behaviour. *Int J Behav Nutr Phys Act*. 2020;17(1):150. doi:10.1186/s12966-020-01040-4
21. Lee SH, Son C, Yeo S, Ha IH. Cross-sectional analysis of self-reported sedentary behaviors and chronic knee pain among South Korean adults over 50 years of age in KNHANES 2013-2015. *BMC Public Health*. 2019;19(1):1375. doi:10.1186/s12889-019-7653-9
22. Dzakpasu FQS, Carver A, Brakenridge CJ, et al. Musculoskeletal pain and sedentary behaviour in occupational and non-occupational settings: a systematic review with meta-analysis. *Int J Behav Nutr Phys Act*. 2021;18(1):159. doi:10.1186/s12966-021-01191-y
23. MedlinePlus. Health Risks of an Inactive Lifestyle. MedlinePlus (National Library of Medicine). Published September 1, 2017. Accessed May 28, 2025. <https://medlineplus.gov/healthrisks/sofainactivelifestyle.html>
24. Soto-Rodríguez FJ, Cabañas EI, Pérez-Mármol JM. Impact of prolonged sitting interruption strategies on shear rate, flow-mediated dilation and blood flow in adults: A systematic review and meta-analysis of randomized cross-over trials. *J Sports Sci*. 2022;40(14):1558-1567. doi:10.1080/02640414.2022.2091347
25. Sackner MA, Patel S, Adams JA. Changes of blood pressure following initiation of physical inactivity and after external addition of pulses to circulation. *Eur J Appl Physiol*. 2019;119(1):201-211. doi:10.1007/s00421-018-4016-7
26. Hooker SP, Diaz KM, Blair SN, et al. Association of accelerometer-measured sedentary time and physical activity with risk of stroke among US adults. *JAMA Netw Open*. 2022;5(6):e2215385. doi:10.1001/jamanetworkopen.2022.15385
27. Peter-Marske KM, Evenson KR, Moore CC, et al. Association of accelerometer-measured physical activity and sedentary behavior with incident cardiovascular disease, myocardial infarction, and ischemic stroke: The Women's Health Study. *J Am Heart Assoc*. 2023;12(7):e028180. doi:10.1161/JAHA.122.028180
28. Ajufo E, Kany S, Rämö JT, et al. Accelerometer-measured sedentary behavior AND risk OF future cardiovascular disease. *J Am Coll Cardiol*. 2025;85(5):473-486. doi:10.1016/j.jacc.2024.10.065
29. Burini RC, Anderson E, Durstine JL, Carson JA. Inflammation, physical activity, and chronic disease: An evolutionary perspective. *Sports Med Health Sci*. 2020;2(1):1-6. doi:10.1016/j.smhs.2020.03.004
30. Pinto AJ, Bergouignan A, Dempsey PC, et al. Physiology of sedentary behavior. *Physiol Rev*. 2023;103(4):2561-2622. doi:10.1152/physrev.00022.2022
31. Damiot A, Pinto AJ, Turner JE, Gualano B. Immunological implications of physical inactivity among older adults during the covid-19 pandemic. *Gerontology*. Published online June 25, 2020:1-8. doi:10.1159/000509216

32. Adil S, Gordon M, Hathagoda W, Kuruppu C, Benninga MA, Rajindrajith S. Impact of physical inactivity and sedentary behaviour on functional constipation in children and adolescents: a systematic review. *BMJ Paediatr Open*. 2024;8(1):e003069. doi:10.1136/bmjpo-2024-003069
33. Pérez-Prieto I, Plaza-Flórida A, Ubago-Guisado E, Ortega FB, Altmäe S. Physical activity, sedentary behavior and microbiome: A systematic review and meta-analysis. *J Sci Med Sport*. 2024;27(11):793-804. doi:10.1016/j.jsams.2024.07.003
34. MedlinePlus. Are you getting too much exercise?. MedlinePlus (National Library of Medicine). Published September 4, 2024. Accessed May 28, 2025. <https://medlineplus.gov/ency/patientinstructions/000807.htm>
35. NIA. Three Types of Exercise Can Improve Your Health and Physical Ability. NIH National Institute on Aging. Published January 14, 2025. Accessed May 28, 2025. <https://www.nia.nih.gov/health/exercise-and-physical-activity/three-types-exercise-can-improve-your-health-and-physical>
36. Daneshmandi H, Choobineh A, Ghaem H, Karimi M. Adverse effects of prolonged sitting behavior on the general health of office workers. *J Lifestyle Med*. 2017;7(2):69-75. doi:10.15280/jlm.2017.7.2.69
37. Zhou L, Deng X, Xu M, et al. The effects of active workstations on reducing work-specific sedentary time in office workers: a network meta-analysis of 23 randomized controlled trials. *Int J Behav Nutr Phys Act*. 2023;20:92. doi:10.1186/s12966-023-01467-5