
Elimination Diet— Bibliography

Scientific/Medical Publications

Features of the Elimination Diet

- Dieterich W, Schuppan D, Schink M, et al. Influence of low FODMAP and gluten-free diets on disease activity and intestinal microbiota in patients with non-celiac gluten sensitivity. Clin Nutr. 2019;38(2):697-707. doi:10.1016/j.clnu.2018.03.017.
- Choghakhori R, Abbasnezhad A, Hasanvand A, Amani R. Inflammatory cytokines and oxidative stress biomarkers in irritable bowel syndrome: association with digestive symptoms and quality of life. Cytokine. 2017;93:34-43. doi:10.1016/j.cyto.2017.05.005.
- Yang J, Deng Y, Chu H, et al. Prevalence and presentation of lactose intolerance and effects on dairy product intake in healthy subjects and patients with irritable bowel syndrome. Clin Gastroenterol Hepatol. 2013;11(3):262-268.e1. doi:10.1016/j.cgh.2012.11.034.
- Gupta D, Ghoshal UC, Misra A, et al. Lactose intolerance in patients with irritable bowel syndrome from northern India: a case-control study. J Gastroenterol Hepatol. 2007;22(12):2261-2265. doi:10.1111/j.1440-1746.2007.04986.x.
- Jianqin S, Leiming X, Lu X, et al. Effects of milk containing only A2 beta casein versus milk containing both A1 and A2 beta casein proteins on gastrointestinal physiology, symptoms of discomfort, and cognitive behavior of people with self-reported intolerance to traditional cows' milk. Nutr J. 2016;15:35. Published 2016 Apr 2. doi:10.1186/s12937-016-0147-z.
- Di Sabatino A, Volta U, Salvatore C, et al. Small amounts of gluten in subjects with suspected nonceliac gluten sensitivity: a randomized, double-blind, placebo-controlled, cross-over trial. Clin Gastroenterol Hepatol. 2015;13(9):1604-12.e3. doi:10.1016/j.cgh.2015.01.029.
- Isasi C, Tejerina E, Fernandez-Puga N, Serrano-Vela JL. Fibromyalgia and chronic fatigue syndrome caused by non-celiac gluten sensitivity. Reumatol Clin. 2015;11(1):56-57. doi:10.1016/j.reuma.2014.06.005.

Scientific/Medical Publications

Features of the Elimination Diet (continued)

- Griausdaite K, Maselis K, Žvirbliene A, et al. Associations between migraine, celiac disease, non-celiac gluten sensitivity and activity of diamine oxidase. *Med Hypotheses*. 2020;142:109738. doi:10.1016/j.mehy.2020.109738.
- Daley CA, Abbott A, Doyle PS, et al. A review of fatty acid profiles and antioxidant content in grass-fed and grain-fed beef. *Nutr J*. 2010;9:10. Published 2010 Mar 10. doi:10.1186/1475-2891-9-10.

Touring Through the Food Plan

- Gwin JA, Maki KC, Leidy HJ. Increased protein consumption during the day from an energy-restricted diet augments satiety but does not reduce daily fat or carbohydrate intake on a free-living test day in overweight women. *J Nutr*. 2017;147(12):2338-2346. doi:10.3945/jn.117.255554.
- Welsh JA, Braun H, Brown N, et al. Production-related contaminants (pesticides, antibiotics and hormones) in organic and conventionally produced milk samples sold in the USA. *Public Health Nutr*. 2019;22(16):2972-2980. doi:10.1017/S136898001900106X.
- Dikariyanto V, Smith L, Francis L, et al. Snacking on whole almonds for 6 weeks improves endothelial function and lowers LDL cholesterol but does not affect liver fat and other cardiometabolic risk factors in healthy adults: the ATTIS study, a randomized controlled trial. *Am J Clin Nutr*. 2020;111(6):1178-1189. doi:10.1093/ajcn/nqaa100.
- Jung H, Chen CO, Blumberg JB, Kwak HK. The effect of almonds on vitamin E status and cardiovascular risk factors in Korean adults: a randomized clinical trial. *Eur J Nutr*. 2018;57(6):2069-2079. doi:10.1007/s00394-017-1480-5.
- Guasch-Ferré M, Hu FB, Martínez-González MA, et al. Olive oil intake and risk of cardiovascular disease and mortality in the PREDIMED Study. *BMC Med*. 2014;12:78. Published 2014 May 13. doi:10.1186/1741-7015-12-78.
- Pintó X, Fanlo-Maresma M, Corbella E, et al. A Mediterranean diet rich in extra-virgin olive oil is associated with a reduced prevalence of nonalcoholic fatty liver disease in older individuals at high cardiovascular risk. *J Nutr*. 2019;149(11):1920-1929. doi:10.1093/jn/nxz147.
- Aslam F, Iqbal S, Nasir M, Anjum AA. White sesame seed oil mitigates blood glucose level, reduces oxidative stress, and improves biomarkers of hepatic and renal function in participants with type 2 diabetes mellitus. *J Am Coll Nutr*. 2019;38(3):235-246. doi:10.1080/07315724.2018.1500183.
- Chen JG, Johnson J, Egner P, et al. Dose-dependent detoxication of the airborne pollutant benzene in a randomized trial of broccoli sprout beverage in Qidong, China. *Am J Clin Nutr*. 2019 Sep 1;110(3):675-684. doi: 10.1093/ajcn/nqz122.

Scientific/Medical Publications

Touring Through the Food Plan (continued)

- Egner PA, Chen JG, Zarth AT, et al. Rapid and sustainable detoxication of airborne pollutants by broccoli sprout beverage: results of a randomized clinical trial in China. *Cancer Prev Res (Phila)*. 2014;7(8):813-823. doi:10.1158/1940-6207.CAPR-14-0103.
- Du C, Smith A, Avalos M, et al. Blueberries improve pain, gait performance, and inflammation in individuals with symptomatic knee osteoarthritis. *Nutrients*. 2019;11(2):290. Published 2019 Jan 29. doi:10.3390/nu11020290.
- Schell J, Scofield RH, Barrett JR, et al. Strawberries improve pain and inflammation in obese adults with radiographic evidence of knee osteoarthritis. *Nutrients*. 2017;9(9):949. Published 2017 Aug 28. doi:10.3390/nu9090949.
- Boespflug EL, Eliassen JC, Dudley JA, et al. Enhanced neural activation with blueberry supplementation in mild cognitive impairment. *Nutr Neurosci*. 2018;21(4):297-305. doi:10.1080/1028415X.2017.1287833.
- Barfoot KL, May G, Lamport DJ, et al. The effects of acute wild blueberry supplementation on the cognition of 7-10-year-old schoolchildren. *Eur J Nutr*. 2019;58(7):2911-2920. doi:10.1007/s00394-018-1843-6.
- Curtis PJ, van der Velpen V, Berends L, et al. Blueberries improve biomarkers of cardiometabolic function in participants with metabolic syndrome-results from a 6-month, double-blind, randomized controlled trial. *Am J Clin Nutr*. 2019;109(6):1535-1545. doi:10.1093/ajcn/nqy380.
- Eady SL, Wallace AJ, Hedderley DI, et al. The effects on immune function and digestive health of consuming the skin and flesh of Zespri® SunGold kiwifruit (*Actinidia chinensis* var. *chinensis* 'Zesy002') in healthy and IBS-constipated individuals. *Nutrients*. 2020;12(5):1453. Published 2020 May 18. doi:10.3390/nu12051453.
- Barati Boldaji R, Akhlaghi M, Sagheb MM, Esmaeilinezhad Z. Pomegranate juice improves cardiometabolic risk factors, biomarkers of oxidative stress and inflammation in hemodialysis patients: a randomized crossover trial. *J Sci Food Agric*. 2020;100(2):846-854. doi:10.1002/jsfa.10096.
- Sohrab G, Roshan H, Ebrahimof S, et al. Effects of pomegranate juice consumption on blood pressure and lipid profile in patients with type 2 diabetes: a single-blind randomized clinical trial. *Clin Nutr ESPEN*. 2019;29:30-35. doi:10.1016/j.clnesp.2018.11.013.
- Kojadinovic MI, Arsic AC, Debeljak-Martacic JD, et al. Consumption of pomegranate juice decreases blood lipid peroxidation and levels of arachidonic acid in women with metabolic syndrome. *J Sci Food Agric*. 2017;97(6):1798-1804. doi:10.1002/jsfa.7977.

Scientific/Medical Publications

Touring Through the Food Plan (continued)

- Bondonno NP, Bondonno CP, Blekkenhorst LC, et al. Flavonoid-rich apple improves endothelial function in individuals at risk for cardiovascular disease: a randomized controlled clinical trial. *Mol Nutr Food Res*. 2018;62(3):10.1002/mnfr.201700674. doi:10.1002/mnfr.201700674.
- Bondonno CP, Yang X, Croft KD, et al. Flavonoid-rich apples and nitrate-rich spinach augment nitric oxide status and improve endothelial function in healthy men and women: a randomized controlled trial. *Free Radic Biol Med*. 2012;52(1):95-102. doi:10.1016/j.freeradbiomed.2011.09.028.
- Roager HM, Vogt JK, Kristensen M, et al. Whole grain-rich diet reduces body weight and systemic low-grade inflammation without inducing major changes of the gut microbiome: a randomised cross-over trial. *Gut*. 2019;68(1):83-93. doi:10.1136/gutjnl-2017-314786.
- Kopf JC, Suhr MJ, Clarke J, et al. Role of whole grains versus fruits and vegetables in reducing subclinical inflammation and promoting gastrointestinal health in individuals affected by overweight and obesity: a randomized controlled trial. *Nutr J*. 2018;17(1):72. Published 2018 Jul 30. doi:10.1186/s12937-018-0381-7.
- Anti M, Pignataro G, Armuzzi A, et al. Water supplementation enhances the effect of high-fiber diet on stool frequency and laxative consumption in adult patients with functional constipation. *Hepatogastroenterology*. 1998;45(21):727-732.
- Jeong JN. Effect of pre-meal water consumption on energy intake and satiety in non-obese young adults. *Clin Nutr Res*. 2018;7(4):291-296. doi:10.7762/cnr.2018.7.4.291.
- Zhang J, Zhang N, He H, Du S, Ma G. Different amounts of water supplementation improved cognitive performance and mood among young adults after 12 h water restriction in Baoding, China: a randomized controlled trial (RCT). *Int J Environ Res Public Health*. 2020;17(21):7792. Published 2020 Oct 24. doi:10.3390/ijerph17217792.
- Zhang N, Du SM, Zhang JF, Ma GS. Effects of dehydration and rehydration on cognitive performance and mood among male college students in Cangzhou, China: a self-controlled trial. *Int J Environ Res Public Health*. 2019;16(11):1891. Published 2019 May 29. doi:10.3390/ijerph16111891.
- Shearer J, Reimer RA, Hittel DS, et al. Caffeine-containing energy shots cause acute impaired glucoregulation in adolescents. *Nutrients*. 2020;12(12):3850. Published 2020 Dec 16. doi:10.3390/nu12123850.
- Beaudoin MS, Allen B, Mazzetti G, et al. Caffeine ingestion impairs insulin sensitivity in a dose-dependent manner in both men and women. *Appl Physiol Nutr Metab*. 2013;38(2):140-147. doi:10.1139/apnm-2012-0201.

Scientific/Medical Publications

Touring Through the Food Plan (continued)

- Hagen K, Thoresen K, Stovner LJ, Zwart JA. High dietary caffeine consumption is associated with a modest increase in headache prevalence: results from the Head-HUNT Study. *J Headache Pain*. 2009;10(3):153-159. doi:10.1007/s10194-009-0114-6.
- Sharifzadeh F, Kashanian M, Koohpayehzadeh J, et al. A comparison between the effects of ginger, pyridoxine (vitamin B6) and placebo for the treatment of the first trimester nausea and vomiting of pregnancy (NVP). *J Matern Fetal Neonatal Med*. 2018;31(19):2509-2514. doi:10.1080/14767058.2017.1344965.
- Sanaati F, Najafi S, Kashaninia Z, Sadeghi M. Effect of ginger and chamomile on nausea and vomiting caused by chemotherapy in Iranian women with breast cancer. *Asian Pac J Cancer Prev*. 2016;17(8):4125-4129.
- Uchio R, Muroyama K, Okuda-Hanafusa C, et al. Hot water extract of *Curcuma longa* L. improves serum inflammatory markers and general health in subjects with overweight or prehypertension/mild hypertension: a randomized, double-blind, placebo-controlled trial. *Nutrients*. 2019;11(8):1822. Published 2019 Aug 7. doi:10.3390/nu11081822.
- Kuptniratsaikul V, Thanakhumtorn S, Chinswangwatanakul P, et al. Efficacy and safety of *Curcuma domestica* extracts in patients with knee osteoarthritis. *J Altern Complement Med*. 2009;15(8):891-897. doi:10.1089/acm.2008.0186.

