
Health Benefits of Spices— Bibliography

Scientific/Medical Publications

- Oh ES, Petersen KS, Kris-Etherton PM, Rogers CJ. Spices in a high-saturated-fat, high-carbohydrate meal reduce postprandial proinflammatory cytokine secretion in men with overweight or obesity: a 3-period, crossover, randomized controlled trial. *J Nutr.* 2020;150(6):1600-1609. doi:10.1093/jn/nxaa063.
- Petersen KS, Rogers CJ, West SG, et al. The effect of culinary doses of spices in a high-saturated fat, high-carbohydrate meal on postprandial lipemia and endothelial function: a randomized, controlled, crossover pilot trial. *Food Funct.* 2020;11(4):3191-3200. doi:10.1039/c9fo02438g.
- Nakayama H, Tsuge N, Sawada H, et al. A single consumption of curry improved postprandial endothelial function in healthy male subjects: a randomized, controlled crossover trial. *Nutr J.* 2014;13:67. doi:10.1186/1475-2891-13-67.
- Lu QY, Rasmussen AM, Yang J, et al. Mixed spices at culinary doses have prebiotic effects in healthy adults: a pilot study. *Nutrients.* 2019;11(6):1425. doi:10.3390/nu11061425.
- Khine WWT, Halder S, De Loi S, Lee YK. A single serving of mixed spices alters gut microflora composition: a dose-response randomised trial. *Sci Rep.* 2021;11(1):11264. doi:10.1038/s41598-021-90453-7.
- Halder S, Lim J, Chia SC, et al. Effects of two doses of curry prepared with mixed spices on postprandial ghrelin and subjective appetite responses—a randomized controlled crossover trial. *Foods.* 2018;7(4):47. doi:10.3390/foods7040047.
- Chopan M, Littenberg B. The association of hot red chili pepper consumption and mortality: a large population-based cohort study. *PLoS One.* 2017;12(1):e0169876. doi:10.1371/journal.pone.0169876.
- Bonaccio M, Di Castelnuovo A, Costanzo S, et al. Chili pepper consumption and mortality in Italian adults. *J Am Coll Cardiol.* 2019;74(25):3139-3149. doi:10.1016/j.jacc.2019.09.068.
- Cheshmeh S, Ghayyem M, Khamooshi F, et al. Green cardamom plus low-calorie diet can decrease the expression of inflammatory genes among obese women with polycystic ovary syndrome: a double-blind randomized clinical trial. *Eat Weight Disord.* 2021;1-10. doi:10.1007/s40519-021-01223-3.

Scientific/Medical Publications

- Kazemi S, Yaghooblou F, Siassi F, et al. Cardamom supplementation improves inflammatory and oxidative stress biomarkers in hyperlipidemic, overweight, and obese pre-diabetic women: a randomized double-blind clinical trial. *J Sci Food Agric.* 2017;97(15):5296-5301. doi:10.1002/jsfa.8414.
- Azimi P, Ghiasvand R, Feizi A, et al. Effects of cinnamon, cardamom, saffron, and ginger consumption on markers of glycemic control, lipid profile, oxidative stress, and inflammation in type 2 diabetes patients. *Rev Diabet Stud.* 2014;11(3-4):258-266. doi:10.1900/RDS.2014.11.258.
- Aghasi M, Koohdani F, Qorbani M, et al. Beneficial effects of green cardamom on serum SIRT1, glycemic indices and triglyceride levels in patients with type 2 diabetes mellitus: a randomized double-blind placebo controlled clinical trial. *J Sci Food Agric.* 2019;99(8):3933-3940. doi:10.1002/jsfa.9617.
- Daneshi-Maskooni M, Keshavarz SA, Qorbani M, et al. Green cardamom supplementation improves serum irisin, glucose indices, and lipid profiles in overweight or obese non-alcoholic fatty liver disease patients: a double-blind randomized placebo-controlled clinical trial. *BMC Complement Altern Med.* 2019;19(1):59. doi:10.1186/s12906-019-2465-0.
- Shishehbor F, Rezaeyan Safar M, Rajaei E, Haghighizadeh MH. Cinnamon consumption improves clinical symptoms and inflammatory markers in women with rheumatoid arthritis. *J Am Coll Nutr.* 2018;1-6. doi:10.1080/07315724.2018.1460733.
- Shirzad F, Morovatdar N, Rezaee R, et al. Cinnamon effects on blood pressure and metabolic profile: a double-blind, randomized, placebo-controlled trial in patients with stage 1 hypertension. *Avicenna J Phytomed.* 2021;11(1):91-100..
- Borzoei A, Rafrat M, Asghari-Jafarabadi M. Cinnamon improves metabolic factors without detectable effects on adiponectin in women with polycystic ovary syndrome. *Asia Pac J Clin Nutr.* 2018;27(3):556-563. doi:10.6133/apjcn.062017.13.
- Hajimonfarednejad M, Nimrouzi M, Heydari M, et al. Insulin resistance improvement by cinnamon powder in polycystic ovary syndrome: a randomized double-blind placebo controlled clinical trial. *Phytother Res.* 2018;32(2):276-283. doi:10.1002/ptr.5970.
- Gupta Jain S, Puri S, Misra A, et al. Effect of oral cinnamon intervention on metabolic profile and body composition of Asian Indians with metabolic syndrome: a randomized double-blind control trial. *Lipids Health Dis.* 2017;16(1):113. doi:10.1186/s12944-017-0504-8.
- Heshmati J, Sepidarkish M, Morvaridzadeh M, et al. The effect of cinnamon supplementation on glycemic control in women with polycystic ovary syndrome: a systematic review and meta-analysis. *J Food Biochem.* 2021;45(1):e13543. doi:10.1111/jfbc.13543.

Scientific/Medical Publications

- Mirfeizi M, Mehdizadeh Tourzani Z, Mirfeizi SZ, et al. Controlling type 2 diabetes mellitus with herbal medicines: a triple-blind randomized clinical trial of efficacy and safety. *J Diabetes*. 2016;8(5):647-656. doi:10.1111/1753-0407.12342.
- Farhangi MA, Tajmiri S. The effects of powdered black cumin seeds on markers of oxidative stress, intracellular adhesion molecule (ICAM)-1 and vascular cell adhesion molecule (VCAM)-1 in patients with Hashimoto's thyroiditis. *Clin Nutr ESPEN*. 2020;37:207-212. doi:10.1016/j.clnesp.2020.02.015.
- Zare R, Heshmati F, Fallahzadeh H, Nadjarzadeh A. Effect of cumin powder on body composition and lipid profile in overweight and obese women. *Complement Ther Clin Pract*. 2014;20(4):297-301. doi:10.1016/j.ctcp.2014.10.001.
- Farhangi MA, Dehghan P, Tajmiri S. Powdered black cumin seeds strongly improves serum lipids, atherogenic index of plasma and modulates anthropometric features in patients with Hashimoto's thyroiditis. *Lipids Health Dis*. 2018;17(1):59. doi:10.1186/s12944-018-0704-x.
- Taghizadeh M, Memarzadeh MR, Asemi Z, Esmailzadeh A. Effect of the cumin cyminum L. intake on weight loss, metabolic profiles and biomarkers of oxidative stress in overweight subjects: a randomized double-blind placebo-controlled clinical trial. *Ann Nutr Metab*. 2015;66(2-3):117-124. doi:10.1159/000373896.
- Portincasa P, Bonfrate L, Scribano ML, et al. Curcumin and fennel essential oil improve symptoms and quality of life in patients with irritable bowel syndrome. *J Gastrointest Liver Dis*. 2016;25(2):151-157. doi:10.15403/jgld.2014.1121.252.ccm.
- Rahimikian F, Rahimi R, Golzareh P, et al. Effect of *Foeniculum vulgare* Mill. (fennel) on menopausal symptoms in postmenopausal women: a randomized, triple-blind, placebo-controlled trial. *Menopause*. 2017;24(9):1017-1021. doi:10.1097/GME.0000000000000881.
- Zeb F, Safdar M, Fatima S, et al. Supplementation of garlic and coriander seed powder: impact on body mass index, lipid profile and blood pressure of hyperlipidemic patients. *Pak J Pharm Sci*. 2018;31(5):1935-1941.
- Zare E, Alirezaei A, Bakhtiyari M, Mansouri A. Evaluating the effect of garlic extract on serum inflammatory markers of peritoneal dialysis patients: a randomized double-blind clinical trial study. *BMC Nephrol*. 2019;20(1):26. doi:10.1186/s12882-019-1204-6.
- Szulinska M, Kregielska-Narozna M, Swiatek J, et al. Garlic extract favorably modifies markers of endothelial function in obese patients -randomized double blind placebo-controlled nutritional intervention. *Biomed Pharmacother*. 2018;102:792-797. doi:10.1016/j.biopha.2018.03.131.

Scientific/Medical Publications

- Ahmadian F, Mozaffari-Khosravi H, Azaraein MH, et al. The effect of consumption of garlic tablet on proteins oxidation biomarkers in postmenopausal osteoporotic women: a randomized clinical trial. *Electron Physician*. 2017;9(11):5670-5675. doi:10.19082/5670.
- Aalami-Harandi R, Karamali M, Asemi Z. The favorable effects of garlic intake on metabolic profiles, hs-CRP, biomarkers of oxidative stress and pregnancy outcomes in pregnant women at risk for pre-eclampsia: randomized, double-blind, placebo-controlled trial. *J Matern Fetal Neonatal Med*. 2015;28(17):2020-2027. doi:10.3109/14767058.2014.977248.
- Shabani E, Sayemiri K, Mohammadpour M. The effect of garlic on lipid profile and glucose parameters in diabetic patients: a systematic review and meta-analysis. *Prim Care Diabetes*. 2019;13(1):28-42. doi:10.1016/j.pcd.2018.07.007.
- Sangouni AA, Mohammad Hosseini Azar MR, Alizadeh M. Effects of garlic powder supplementation on insulin resistance, oxidative stress, and body composition in patients with non-alcoholic fatty liver disease: a randomized controlled clinical trial. *Complement Ther Med*. 2020;51:102428. doi:10.1016/j.ctim.2020.102428.
- Dehghani S, Alipoor E, Salimzadeh A, et al. The effect of a garlic supplement on the pro-inflammatory adipocytokines, resistin and tumor necrosis factor-alpha, and on pain severity, in overweight or obese women with knee osteoarthritis. *Phytomedicine*. 2018;48:70-75. doi:10.1016/j.phymed.2018.04.060.
- Morvaridzadeh M, Sadeghi E, Agah S, et al. Effect of ginger (*Zingiber officinale*) supplementation on oxidative stress parameters: a systematic review and meta-analysis. *J Food Biochem*. 2021;45(2):e13612. doi:10.1111/jfbc.13612.
- Kamali A, Beigi S, Shokrpour M, Pazuki S. The efficacy of ginger and doxedetomidine in reducing postoperative nausea and vomiting in patients undergoing abdominal hysterectomy. *Altern Ther Health Med*. 2020;26(2):28-33.
- Sharifzadeh F, Kashanian M, Koohpayehzadeh J, et al. A comparison between the effects of ginger, pyridoxine (vitamin B6) and placebo for the treatment of the first trimester nausea and vomiting of pregnancy (NVP). *J Matern Fetal Neonatal Med*. 2018;31(19):2509-2514. doi:10.1080/14767058.2017.1344965.
- Marx W, McCarthy AL, Ried K, et al. The effect of a standardized ginger extract on chemotherapy-induced nausea-related quality of life in patients undergoing moderately or highly emetogenic chemotherapy: a double blind, randomized, placebo controlled trial. *Nutrients*. 2017;9(8):867. doi:10.3390/nu9080867.
- Rafie R, Hosseini SA, Hajiani E, et al. Effect of ginger powder supplementation in patients with non-alcoholic fatty liver disease: a randomized clinical trial. *Clin Exp Gastroenterol*. 2020;13:35-45. doi:10.2147/CEG.S234698.

Scientific/Medical Publications

- Daily JW, Zhang X, Kim DS, Park S. Efficacy of ginger for alleviating the symptoms of primary dysmenorrhea: a systematic review and meta-analysis of randomized clinical trials. *Pain Med.* 2015;16(12):2243-2255. doi:10.1111/pme.12853.
- Shirvani MA, Motahari-Tabari N, Alipour A. The effect of mefenamic acid and ginger on pain relief in primary dysmenorrhea: a randomized clinical trial. *Arch Gynecol Obstet.* 2015;291(6):1277-1281. doi:10.1007/s00404-014-3548-2.
- Kashefi F, Khajehei M, Tabatabaeichehr M, Alavinia M, Asili J. Comparison of the effect of ginger and zinc sulfate on primary dysmenorrhea: a placebo-controlled randomized trial. *Pain Manag Nurs.* 2014;15(4):826-833. doi:10.1016/j.pmn.2013.09.001.
- Black CD, Herring MP, Hurley DJ, O'Connor PJ. Ginger (*Zingiber officinale*) reduces muscle pain caused by eccentric exercise. *J Pain.* 2010;11(9):894-903. doi:10.1016/j.jpain.2009.12.013.
- Farokhnia M, Shafiee Sabet M, Iranpour N, et al. Comparing the efficacy and safety of *Crocus sativus* L. with memantine in patients with moderate to severe Alzheimer's disease: a double-blind randomized clinical trial. *Hum Psychopharmacol.* 2014;29(4):351-359. doi:10.1002/hup.2412.
- Akhondzadeh S, Sabet MS, Harirchian MH, et al. Saffron in the treatment of patients with mild to moderate Alzheimer's disease: a 16-week, randomized and placebo-controlled trial. *J Clin Pharm Ther.* 2010;35(5):581-588. doi:10.1111/j.1365-2710.2009.01133.x.
- Karimi-Nazari E, Nadjarzadeh A, Masoumi R, et al. Effect of saffron (*Crocus sativus* L.) on lipid profile, glycemic indices and antioxidant status among overweight/obese prediabetic individuals: a double-blinded, randomized controlled trial. *Clin Nutr ESPEN.* 2019;34:130-136. doi:10.1016/j.clnesp.2019.07.012.
- Panahi Y, Hosseini MS, Khalili N, et al. Antioxidant and anti-inflammatory effects of curcuminoid-piperine combination in subjects with metabolic syndrome: a randomized controlled trial and an updated meta-analysis. *Clin Nutr.* 2015;34(6):1101-1108. doi:10.1016/j.clnu.2014.12.019.
- Chandran B, Goel A. A randomized, pilot study to assess the efficacy and safety of curcumin in patients with active rheumatoid arthritis. *Phytother Res.* 2012;26(11):1719-1725. doi:10.1002/ptr.4639.
- Panahi Y, Hosseini MS, Khalili N, et al. Effects of curcumin on serum cytokine concentrations in subjects with metabolic syndrome: a post-hoc analysis of a randomized controlled trial. *Biomed Pharmacother.* 2016;82:578-582. doi:10.1016/j.biopha.2016.05.037.
- Lee MS, Wahlqvist ML, Chou YC, et al. Turmeric improves post-prandial working memory in pre-diabetes independent of insulin. *Asia Pac J Clin Nutr.* 2014;23(4):581-591. doi:10.6133/apjcn.2014.23.4.24.

Scientific/Medical Publications

- Lopresti AL, Drummond PD. Efficacy of curcumin, and a saffron/curcumin combination for the treatment of major depression: a randomised, double-blind, placebo-controlled study. *J Affect Disord.* 2017;207:188-196. doi:10.1016/j.jad.2016.09.047.
- Atefi M, Darand M, Entezari MH, et al. A systematic review of the clinical use of curcumin for the management of gastrointestinal diseases. *Adv Exp Med Biol.* 2021;1291:295-326. doi:10.1007/978-3-030-56153-6_18.
- Adab Z, Eghtesadi S, Vafa MR, et al. Effect of turmeric on glycemic status, lipid profile, hs-CRP, and total antioxidant capacity in hyperlipidemic type 2 diabetes mellitus patients. *Phytother Res.* 2019;33(4):1173-1181. doi:10.1002/ptr.6312.
- Ghaffari A, Rafrat M, Navekar R, et al. Turmeric and chicory seed have beneficial effects on obesity markers and lipid profile in non-alcoholic fatty liver disease (NAFLD). *Int J Vitam Nutr Res.* 2019;89(5-6):293-302. doi:10.1024/0300-9831/a000568.
- Navekar R, Rafrat M, Ghaffari A, et al. Turmeric supplementation improves serum glucose indices and leptin levels in patients with nonalcoholic fatty liver diseases. *J Am Coll Nutr.* 2017;36(4):261-267. doi:10.1080/07315724.2016.1267597.
- Rahmani S, Asgary S, Askari G, et al. Treatment of non-alcoholic fatty liver disease with curcumin: a randomized placebo-controlled trial. *Phytother Res.* 2016;30(9):1540-1548. doi:10.1002/ptr.5659.

