
The Role of the Liver in Detoxification

Every day, your body is exposed to harmful chemicals. Your liver,¹ kidneys,² large intestine,³ lymphatic system,⁴ and sweat glands^{5,6} work together to reduce the buildup of these chemicals. This process is called detoxification, or detox. Of these organ systems, the liver plays one of the largest roles. The liver performs hundreds of functions and is needed to process nutrients and hormones,^{7,8} as well as remove waste products created as the result of normal body functions.⁹

The liver also aids in breaking down and removing external toxins¹⁰ (naturally occurring) and toxicants¹¹ (man-made):

- Farm chemicals, air pollutants, and chemicals from personal care products
- Both prescription and over-the-counter medications
- Food additives, colorings, flavor enhancers, preservatives, and artificial sweeteners
- Alcohols
- Volatile organic compounds such as those found in fragrances and air fresheners

Phases of Detoxification in the Liver

Phase I Liver detoxification is the first line of defense against toxins/toxicants. Phase I functions as a result of a group of enzymes known as the Cytochrome P450 family.¹² These enzymes help neutralize substances such as caffeine and alcohol, and also help convert chemicals into forms that can be more easily removed from the body. If these toxic intermediates are allowed to build up, however, they can damage DNA and proteins. Phase II Liver detoxification's role is to help neutralize these intermediates and transform them to compounds that can be removed by the body through a process known as conjugation.¹³

Supporting the Work of the Liver

In functional medicine, diet is used to support the work of the liver in its detox processes. Various nutrients are required to fuel detox pathways.¹⁴ A shortage or deficiency of any one of them could mean an increased body burden, or buildup of chemicals. Additionally, specific foods can support metabolic processes of changing toxic chemicals and waste products to less harmful forms and assist in their elimination.¹⁵

The dietary plans to support detox generally focus on adding in natural and whole foods to support, regulate, start, or stop various processes related to Phase I and Phase II detoxification in the liver.¹⁶ For people with genetic variability in the Cytochrome P450 system, enzyme activity may be impaired or reduced.^{17,18} In these cases, detox food plans that improve Phase I metabolism and Phase II conjugation are recommended.

REFERENCES

1. Trefts E, Gannon M, Wasserman DH. The liver. *Curr Biol*. 2017;27(21):R1147-R1151. doi:10.1016/j.cub.2017.09.019
2. Robson L. The kidney – an organ of critical importance in physiology. *J Physiol*. 2014;592(18):3953-3954. doi:10.1113/jphysiol.2014.279216
3. Shimizu M. Modulation of intestinal functions by dietary substances: an effective approach to health promotion. *J Tradit Complement Med*. 2012;2(2):81-83. doi:10.1016/S2225-4110(16)30080-3
4. Cueni LN, Detmar M. The lymphatic system in health and disease. *Lymphat Res Biol*. 2008;6(3-4):109-122. doi:10.1089/lrb.2008.1008
5. Baker LB. Physiology of sweat gland function: The roles of sweating and sweat composition in human health. *Temperature*. 2019;6(3):211-259. doi:10.1080/23328940.2019.1632145
6. Kuan WH, Chen YL, Liu CL. Excretion of ni, pb, cu, as, and hg in sweat under two sweating conditions. *Int J Environ Res Public Health*. 2022;19(7):4323. doi:10.3390/ijerph19074323
7. Rhyu J, Yu R. Newly discovered endocrine functions of the liver. *World J Hepatol*. 2021;13(11):1611-1628. doi:10.4254/wjh.v13.i11.1611
8. Kasarinaite A, Sinton M, Saunders PTK, Hay DC. The influence of sex hormones in liver function and disease. *Cells*. 2023;12(12):1604. doi:10.3390/cells12121604
9. Tarasenko TN, McGuire PJ. The liver is a metabolic and immunologic organ: a reconsideration of metabolic decompensation due to infection in inborn errors of metabolism (IEM). *Mol Genet Metab*. 2017;121(4):283-288. doi:10.1016/j.ymgme.2017.06.010
10. Qadri SS, Javaid D, Reyaz A, Ganie SY, Reshi MS. Liver disorders and phytotherapy. *Toxicol Rep*. 2025;14:102047. doi:10.1016/j.toxrep.2025.102047
11. Sears ME, Genuis SJ. Environmental determinants of chronic disease and medical approaches: recognition, avoidance, supportive therapy, and detoxification. *J Environ Public Health*. 2012;2012:1-15. doi:10.1155/2012/356798
12. Hodges RE, Minich DM. Modulation of metabolic detoxification pathways using foods and food-derived components: a scientific review with clinical application. *J Nutr Metab*. 2015;2015:1-23. doi:10.1155/2015/760689
13. Panda C, Komarnytsky S, Fleming MN, et al. Guided metabolic detoxification program supports phase II detoxification enzymes and antioxidant balance in healthy participants. *Nutrients*. 2023;15(9):2209. doi:10.3390/nu15092209
14. Jamioł-Milc D, Gudan A, Kaźmierczak-Siedlecka K, et al. Nutritional support for liver diseases. *Nutrients*. 2023;15(16):3640. doi:10.3390/nu15163640
15. Sang R, Jiang W, Zhang C, Yin R, Ouyang Z, Wei Y. Effect of food components on cytochrome P450 expression and activity. *Hum Nutr Metab*. 2025;40:200304. doi:10.1016/j.hnm.2025.200304
16. Mega A, Marzi L, Kob M, Piccin A, Floreani A. Food and nutrition in the pathogenesis of liver damage. *Nutrients*. 2021;13(4):1326. doi:10.3390/nu13041326
17. Hossam Abdelmonem B, Abdelaal NM, Anwer EKE, et al. Decoding the role of cyp450 enzymes in metabolism and disease: a comprehensive review. *Biomedicines*. 2024;12(7):1467. doi:10.3390/biomedicines12071467
18. Goh LL, Lim CW, Sim WC, Toh LX, Leong KP. Analysis of genetic variation in cyp450 genes for clinical implementation. Ahmad A, ed. *PLOS ONE*. 2017;12(1):e0169233. doi:10.1371/journal.pone.0169233